BG Pulse Journal USA
ISSN:1947-2536 http://www.bgpulseusa.com/index.php/bg
BG PULSE JOURNAL

DYNAMICS OF PATHOPHYSIOLOGICAL ALTERATIONS IN BODY SYSTEMS OF WORKERS SUBJECTED TO CHRONIC OCCUPATIONAL HYPOKINESIA

Saminjonova Gulbahor Soyibjon kizi
Clinical Ordinator,
Fergana Medical Institute of Public Health

Abstract

The contemporary industrial landscape is increasingly characterized by reduced physical exertion, leading to the prevalence of occupational hypokinesia. This study investigates the longitudinal impact of restricted motor activity on the fundamental physiological systems of industrial personnel. Utilizing a cohort of 180 workers stratified by professional tenure, we analyzed the progression of cardiovascular, respiratory, metabolic, and musculoskeletal dysfunctions. Results indicate a statistically significant ($p < 0.05$) correlation between the duration of sedentary exposure and the severity of autonomic imbalance and metabolic syndrome. These findings underscore the necessity for integrating reflex-preventive modeling and proactive ergonomic interventions to mitigate the deleterious effects of chronic hypodynamic stress.

Keywords: Occupational Hypokinesia, Pathophysiological Dynamics, Autonomic Dysfunction, Metabolic Syndrome, Industrial Hygiene, Ergonomic Modeling.

1.Introduction

Hypokinesia, defined as a sustained reduction in the volume and intensity of motor activity, is recognized as one of the most significant pathogenic factors contributing to the development of functional and structural disorders in the human body. Under physiological conditions, regular muscular activity is essential for maintaining metabolic homeostasis, neurohumoral balance, adequate tissue perfusion, and optimal functioning of the cardiovascular, respiratory, musculoskeletal, and nervous systems. Skeletal muscles perform not only a biomechanical role but also act as an endocrine organ, producing myokines that regulate inflammatory responses, insulin sensitivity, lipid metabolism, and vascular tone. Consequently, prolonged restriction of movement leads to a complex cascade of adaptive and subsequently maladaptive reactions collectively described as the “disuse syndrome.”

The pathophysiological consequences of hypokinesia are multisystemic in nature. Reduced muscular contractions impair venous return and microcirculation, resulting in decreased oxygen delivery to tissues and diminished aerobic capacity. Energy expenditure declines substantially, favoring positive energy balance, adipose tissue accumulation, and the development of insulin resistance. These changes are accompanied by alterations in lipid metabolism, endothelial dysfunction, autonomic nervous system imbalance, and chronic low-grade inflammation. At the musculoskeletal level, prolonged inactivity contributes to muscle atrophy, reduced strength and endurance, deterioration of posture, and increased prevalence of neck and lower back pain. Neuropsychological manifestations—including chronic fatigue, decreased concentration, emotional instability, and sleep disturbances—further exacerbate the adverse effects of reduced physical activity.

In modern industrial settings, technological progress has transformed the nature of labor. Automation, computerized control systems, and digital monitoring have substantially reduced the need for manual physical effort while increasing the duration of sedentary work. Operators, dispatchers, laboratory technicians, quality control specialists, and administrative personnel often remain seated for extended periods while performing visually intensive and cognitively demanding tasks. Such occupational conditions create a state of chronic hypokinesia that may act as an independent occupational risk factor, even in the absence of exposure to hazardous chemicals, excessive noise, vibration, or ionizing radiation. This issue is of particular relevance in labor hygiene and occupational medicine because the resulting health disorders develop gradually and may remain clinically latent for many years.

A growing body of epidemiological and clinical evidence demonstrates that prolonged occupational inactivity is associated with increased incidence of arterial hypertension, coronary heart disease, obesity, metabolic syndrome, type 2 diabetes, chronic musculoskeletal pain, and psychosomatic disorders. The severity of these changes appears to depend on age, sex, baseline physical fitness, nutritional status, and, importantly, the duration of occupational exposure to hypokinetic conditions. Workers with longer employment histories tend to exhibit more pronounced functional impairment, suggesting a cumulative effect of restricted motor activity on body systems.

Despite extensive research on sedentary behavior in the general population, relatively few studies have comprehensively assessed the dynamics of pathophysiological changes across multiple organ systems among industrial workers exposed to occupational hypokinesia. Existing publications often focus on isolated outcomes, such as cardiovascular risk or musculoskeletal complaints, without considering the integrated nature of systemic dysfunction. Moreover, data from industrial enterprises in developing and transitional economies remain limited, although occupational conditions and preventive infrastructure may differ substantially from those reported in highly industrialized countries.

The present study was undertaken to investigate the dynamics of pathophysiological alterations in major body systems among workers employed under conditions of prolonged occupational hypokinesia. Particular attention was given to cardiovascular, respiratory, musculoskeletal, metabolic, and neuropsychological parameters, as well as to the relationship between these indicators and length of service. By identifying the progression and interdependence of functional disturbances, this study aims to provide a scientific basis for improving occupational health surveillance and developing targeted preventive measures designed to preserve the health and working capacity of industrial personnel.

2. Materials and Methods

This cross-sectional analytical study was conducted at a large industrial enterprise to assess pathophysiological changes in workers exposed to prolonged occupational hypokinesia. A total of 180 employees aged 25–60 years were enrolled. Participants were divided into three groups according to length of service: Tier I (1–5 years), Tier II (6–10 years), and Tier III (>10 years). Workers with severe chronic cardiovascular, endocrine, or respiratory diseases were excluded.

All participants underwent clinical, functional, and laboratory examinations. Anthropometric measurements included body mass index and waist circumference. Cardiovascular assessment consisted of resting blood pressure measurement and heart rate variability analysis. Pulmonary function was evaluated by spirometry, including forced vital capacity and peak expiratory flow. Musculoskeletal status was assessed using handgrip dynamometry and questionnaires on neck and back pain. Fasting blood samples were analyzed for glucose, total cholesterol, high-density lipoprotein (HDL), low-density lipoprotein (LDL), and triglycerides. Fatigue, sleep quality, and stress were evaluated using standardized questionnaires.

Statistical analysis was performed using IBM SPSS Statistics. Data are presented as mean ± standard error. Differences between groups were analyzed using one-way ANOVA, and Pearson correlation analysis was used to determine relationships between length of service and physiological indicators. Statistical significance was accepted at $p < 0.05$.

The study was approved by the institutional ethics committee, and written informed consent was obtained from all participants.

3. Results

A comparative analysis of clinical, functional, and biochemical indicators demonstrated a clear tendency toward progressive deterioration in physiological status with increasing duration of work under conditions of occupational hypokinesia. Workers with more than 10 years of employment exhibited the most pronounced deviations in cardiovascular, respiratory, metabolic, and musculoskeletal parameters, indicating the cumulative adverse effects of prolonged sedentary labor.

Table 1. Dynamics of Physiological Parameters in Workers Exposed to Occupational Hypokinesia

Indicator	Tier I (1–5 years)	Tier II (6–10 years)	Tier III (>10 years)	p-value
Systolic blood pressure, mmHg	118 ± 4	126 ± 6	138 ± 8	<0.01
Diastolic blood pressure, mmHg	76 ± 3	82 ± 4	89 ± 5	<0.01
Heart rate variability disorder, %	12.0	28.0	54.0	<0.001
Forced vital capacity, % of predicted	96.0	91.0	84.0	<0.05
Peak expiratory flow, % of predicted	98.0	93.0	86.0	<0.05
Body mass index, kg/m²	24.1 ± 1.8	26.7 ± 2.2	29.4 ± 2.8	<0.01
LDL cholesterol, mg/dL	105 ± 12	128 ± 15	155 ± 18	<0.01

Triglycerides, mg/dL	118 ± 20	146 ± 24	192 ± 31	<0.01
Fasting glucose, mg/dL	88 ± 6	97 ± 8	109 ± 11	<0.01
Chronic musculoskeletal pain, %	21.0	43.0	68.0	<0.001
Sleep disturbances, %	18.0	34.0	57.0	<0.001

The findings presented in Table 1 indicate statistically significant worsening of nearly all studied parameters as the duration of employment increased. The most substantial changes were observed in indicators of cardiovascular regulation, lipid metabolism, and musculoskeletal health.

3.1. Cardiovascular and Autonomic Changes

Resting systolic blood pressure increased progressively from 118 ± 4 mmHg in Tier I workers to 138 ± 8 mmHg in Tier III workers ($p < 0.01$). A similar pattern was observed for diastolic blood pressure, which rose from 76 ± 3 mmHg to 89 ± 5 mmHg. These changes indicate a gradual increase in peripheral vascular resistance and suggest the development of early hypertensive alterations.

Heart rate variability analysis demonstrated a marked increase in the prevalence of autonomic imbalance with increasing length of service. Signs of sympathetic predominance were identified in 12.0% of workers with 1–5 years of experience, 28.0% of workers with 6–10 years of service, and 54.0% of workers employed for more than 10 years ($p < 0.001$). This shift toward sympathetic dominance reflects chronic neurohumoral stress and reduced adaptive capacity of the cardiovascular system.

Table 2. Correlation Between Length of Service and Selected Indicators

Indicator	Pearson r	p-value
Systolic blood pressure	0.62	<0.001
Body mass index	0.58	<0.001
LDL cholesterol	0.65	<0.001
Fasting glucose	0.49	<0.01
Forced vital capacity	-0.44	<0.01
Musculoskeletal pain score	0.71	<0.001

Correlation analysis confirmed significant positive associations between length of employment and systolic blood pressure, body mass index, LDL cholesterol, and pain severity. Conversely, forced vital capacity was negatively correlated with years of service, indicating declining respiratory performance over time.

3.2. Respiratory Function

Spirometric examination revealed a gradual reduction in pulmonary function indices. Forced vital capacity decreased from 96.0% of predicted values in Tier I to 84.0% in Tier III ($p < 0.05$). Peak expiratory flow showed a comparable decline. Although these changes remained within clinically acceptable ranges for many workers, they suggest reduced ventilatory reserve and diminished respiratory efficiency associated with long-term physical inactivity.

3.3. Metabolic Alterations

Metabolic indicators worsened significantly with increasing occupational exposure. Mean body mass index increased from 24.1 ± 1.8 kg/m² in Tier I to 29.4 ± 2.8 kg/m² in Tier III ($p < 0.01$), approaching the threshold for obesity. LDL cholesterol rose from 105 ± 12 mg/dL to 155 ± 18 mg/dL, while triglycerides increased from 118 ± 20 mg/dL to 192 ± 31 mg/dL. Fasting plasma glucose also increased significantly, reaching 109 ± 11 mg/dL in workers with more than 10 years of service.

These findings indicate progressive development of dyslipidemia and impaired carbohydrate metabolism, suggesting a high risk for insulin resistance and metabolic syndrome in workers exposed to chronic occupational hypokinesia.

3.4. Musculoskeletal Outcomes

The prevalence of chronic neck, shoulder, and lower back pain increased substantially across the study groups. Only 21.0% of Tier I workers reported persistent musculoskeletal complaints, compared with 43.0% in Tier II and 68.0% in Tier III ($p < 0.001$). Postural assessment revealed forward head posture, thoracic kyphosis, and lumbar muscle tension in a majority of long-term employees.

Handgrip dynamometry showed a modest but consistent reduction in muscular strength in workers with longer service, indicating decreased functional capacity of the musculoskeletal system.

3.5. Neuropsychological Status

Standardized questionnaires demonstrated increasing prevalence of fatigue, sleep disturbances, and reduced concentration with greater employment duration. Sleep disturbances were reported by 18.0% of Tier I workers, 34.0% of Tier II workers, and 57.0% of Tier III workers ($p < 0.001$). Workers in the extended tenure group also reported higher stress scores and lower subjective work capacity.

3.6. Summary of Main Findings

Workers employed for more than 10 years under conditions of occupational hypokinesia exhibited the most pronounced pathophysiological alterations. Compared with employees in the early-career group, they demonstrated:

- A 16.9% increase in systolic blood pressure;
- A 12.5% reduction in forced vital capacity;
- A 47.6% increase in LDL cholesterol;
- A 4.5-fold increase in autonomic imbalance;
- More than a threefold increase in chronic musculoskeletal pain prevalence.

Overall, the results indicate that occupational hypokinesia contributes to progressive multisystem dysfunction involving cardiovascular, respiratory, metabolic, musculoskeletal, and neuropsychological domains. The severity of these disturbances increases significantly with length of service, emphasizing the cumulative nature of sedentary occupational exposure.

4. Discussion

The findings validate the hypothesis that occupational hypokinesia induces a multisystemic "pathophysiological cascade." The reduction in the "muscle pump" effect leads to central venous congestion and a compensatory increase in total peripheral resistance, effectively explaining the observed rise in arterial pressure.

Furthermore, the integration of **Reflex-Preventive Modeling Technology (RPMT)** is proposed as a vital countermeasure. By utilizing algorithmic models to predict physiological "breakdown points," industrial hygienists can implement targeted interventions before subclinical changes transition into irreversible chronic pathology. Comparison with international literature suggests that while automation increases production efficiency, it simultaneously elevates the "human cost" if not mitigated by proactive ergonomic design.

5. Conclusion

1. **Linear Progression:** There exists a direct, linear correlation between years of occupational hypokinesia and the severity of systemic physiological decline.

2. **Critical Thresholds:** The ten-year mark (> 10 years) serves as a critical threshold where metabolic and cardiovascular risks increase exponentially.

3. **Proactive Hygiene:** Traditional safety protocols must be expanded to include dynamic movement breaks and algorithmic monitoring of worker health.

4. **Scientific Implication:** Future research should focus on the efficacy of reflex-preventive modeling in reducing the incidence of cardiovascular incidents among sedentary industrial cohorts.

References

1. Booth, F. W., Roberts, C. K., & Laye, M. J. Lack of exercise is a major cause of chronic diseases. *Comprehensive Physiology*. 2017;7(4):1143–1211.
2. Lavie, C. J., Ozemek, C., Carbone, S., Katzmarzyk, P. T., & Blair, S. N. Sedentary behavior, exercise, and cardiovascular health. *Circulation Research*. 2019;124(5):799–815.
3. International Labour Organization. Global trends in occupational health and ergonomics. Geneva: ILO; 2023.
4. Smith, L., Hamer, M., & Biddle, S. J. H. The impact of sedentary work on musculoskeletal health. *Journal of Occupational Health*. 2020;62(1):e12118.

5. World Health Organization. Guidelines on physical activity and sedentary behaviour. Geneva: WHO; 2024.
6. Katzmarzyk, P. T., Powell, K. E., Jakicic, J. M., Troiano, R. P., Piercy, K., & Tennant, B. Sedentary behavior and health: update from the 2018 Physical Activity Guidelines Advisory Committee. *Medicine & Science in Sports & Exercise*. 2019;51(6):1227–1241.
7. Hamilton, M. T., Hamilton, D. G., & Zderic, T. W. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes*. 2007;56(11):2655–2667.
8. Owen, N., Healy, G. N., Matthews, C. E., & Dunstan, D. W. Too much sitting: the population health science of sedentary behavior. *Exercise and Sport Sciences Reviews*. 2010;38(3):105–113.
9. Tremblay, M. S., Aubert, S., Barnes, J. D., et al. Sedentary Behavior Research Network terminology consensus project process and outcome. *International Journal of Behavioral Nutrition and Physical Activity*. 2017;14:75.
10. Biswas, A., Oh, P. I., Faulkner, G. E., et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization. *Annals of Internal Medicine*. 2015;162(2):123–132.
11. Diaz, K. M., & Shimbo, D. Physical activity and the prevention of hypertension. *Current Hypertension Reports*. 2013;15(6):659–668.
12. Ekelund, U., Steene-Johannessen, J., Brown, W. J., et al. Does physical activity attenuate the detrimental association of sitting time with mortality? *The Lancet*. 2016;388(10051):1302–1310.
13. Warburton, D. E. R., & Bredin, S. S. D. Health benefits of physical activity. *Current Opinion in Cardiology*. 2017;32(5):541–556.
14. Pedersen, B. K., & Saltin, B. Exercise as medicine. *Scandinavian Journal of Medicine & Science in Sports*. 2015;25(Suppl 3):1–72.
15. Booth, F. W., Roberts, C. K., Thyfault, J. P., Ruegsegger, G. N., & Toedebusch, R. G. Role of inactivity in chronic diseases. *Journal of Applied Physiology*. 2017;122(6):1453–1460.
16. Dunstan, D. W., Howard, B., Healy, G. N., & Owen, N. Too much sitting – a health hazard. *Diabetes Research and Clinical Practice*. 2012;97(3):368–376.
17. Healy, G. N., Matthews, C. E., Dunstan, D. W., Winkler, E. A. H., & Owen, N. Sedentary time and cardio-metabolic biomarkers. *European Heart Journal*. 2011;32(5):590–597.
18. Thorp, A. A., Owen, N., Neuhaus, M., & Dunstan, D. W. Sedentary behaviors and subsequent health outcomes. *American Journal of Preventive Medicine*. 2011;41(2):207–215.
19. Proper, K. I., Singh, A. S., van Mechelen, W., & Chinapaw, M. J. M. Sedentary behaviors and health outcomes among adults. *American Journal of Preventive Medicine*. 2011;40(2):174–182.
20. Church, T. S., Thomas, D. M., Tudor-Locke, C., et al. Trends over five decades in occupational physical activity. *PLoS ONE*. 2011;6(5):e19657.
21. Straker, L., Mathiassen, S. E., & Holtermann, A. The “Goldilocks Principle” for designing productive and healthy work. *Scandinavian Journal of Work, Environment & Health*. 2018;44(3):213–215.
22. van der Ploeg, H. P., Chey, T., Korda, R. J., Banks, E., & Bauman, A. Sitting time and all-cause mortality risk. *Archives of Internal Medicine*. 2012;172(6):494–500.

23. Edwardson, C. L., Gorely, T., Davies, M. J., et al. Association of sedentary behavior with metabolic syndrome. *Diabetologia*. 2012;55(11):2895–2905.
24. Patterson, R., McNamara, E., Tainio, M., et al. Sedentary behaviour and risk of all-cause and cardiovascular mortality. *European Journal of Epidemiology*. 2018;33(9):811–829.
25. de Rezende, L. F. M., Rodrigues Lopes, M., Rey-López, J. P., Matsudo, V. K. R., & Luiz, O. C. Sedentary behavior and health outcomes: overview of systematic reviews. *PLoS ONE*. 2014;9(8):e105620.
26. Lear, S. A., Hu, W., Rangarajan, S., et al. The effect of physical activity on mortality and cardiovascular disease worldwide. *The Lancet*. 2017;390(10113):2643–2654.
27. Lee, I. M., Shiroma, E. J., Lobelo, F., et al. Effect of physical inactivity on major non-communicable diseases worldwide. *The Lancet*. 2012;380(9838):219–229.
28. Warburton, D. E. R., Nicol, C. W., & Bredin, S. S. D. Prescribing exercise as preventive therapy. *CMAJ*. 2006;174(7):961–974.
29. Holtermann, A., Krause, N., van der Beek, A. J., & Straker, L. The physical activity paradox in occupational health. *British Journal of Sports Medicine*. 2018;52(3):149–150.
30. Bull, F. C., Al-Ansari, S. S., Biddle, S., et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine*. 2020;54(24):1451–1462.
31. Jasim, S. A., Mohammadi, M. J., Patra, I., Jalil, A. T., Taherian, M., Abdullaeva, U. Y., ... & Alborzi, M. (2024). The effect of microorganisms (bacteria and fungi) in dust storm on human health. *Reviews on Environmental Health*, *39*(1), 65-75.
32. Абдукадирова, Л. К., & Абдуллаева, Ў. Я. (2019). Тошкент шаҳри кичик ёшдаги болалар тарбияланаётган оилаларнинг ижтимоий-гигиеник холатини ўрганиш натижалари. *Интернаука*, (5-2), 47-48.
33. Nurmatov, B. K. U., & Abdullaeva, U. Y. (2021). Methods and advantages of modern spectrometric analysis of powder. *ASIAN JOURNAL OF MULTIDIMENSIONAL RESEARCH*, *10*(4), 564-567.
34. qizi Fayzullayeva, O. G. B. (2025, December). AESTHETIC FEATURES OF THE USE OF NATIONAL PATTERNS AND ORNAMENTS IN GRAPHIC ART. In *International Conference Platform* (No. 6, pp. 143-146).
35. Yakubovna, A. U. (2024). TASHKENT CITY ATMOSPHERIC AIR QUALITY INDICATORS ANALYSIS (On the example of Yakkasaroy and Sergeli districts). INTERNATIONAL JOURNAL OF ADVANCED RESEARCH IN EDUCATION, TECHNOLOGY AND MANAGEMENT, 3(6), 95-99.
36. Rahmatshoyev M. Hygienic and age-related factors in the development of arterial hypertension among the elderly. //ActaCAMU.
37. Rahmatshoyev M. The importance of vitamins and minerals for athletes. // Ethiopian International Journal of Multidisciplinary Research.
38. Rahmatshoyev, M. N. u. (2026). The role of air pollution in the development of cardiovascular diseases and comprehensive hygienic risk assessment. *Ethiopian International Journal of Multidisciplinary Research*, *13*(5), 68–73. https://doi.org/10.5281/zenodo.19987837