BG Pulse Journal USA
ISSN:1947-2536 http://www.bgpulseusa.com/index.php/bg
BG PULSE JOURNAL

Obesity and Insulin Resistance: Mechanisms of Development and Their Interrelationship

Seejap Ali Khan
Medical researcher
doi:https://doi.org/10.5281/zenodo.23757342

Abstract: Obesity and insulin resistance are closely associated metabolic conditions that contribute to the development of type 2 diabetes and other chronic diseases. Their relationship involves more than an increase in total body fat. Adipose tissue dysfunction, abnormal lipid distribution, inflammatory signaling, and impaired insulin action in skeletal muscle and the liver are important contributing mechanisms. However, insulin resistance varies considerably among individuals, and obesity does not inevitably lead to diabetes. This narrative overview examines the biological relationship between obesity and insulin resistance and discusses its implications for clinical assessment, prevention, and management.

Keywords: obesity, insulin resistance, adipose tissue, inflammation, ectopic fat, metabolic dysfunction, type 2 diabetes.

Introduction

Obesity is an important risk factor for insulin resistance, a condition in which tissues respond inadequately to insulin. However, these conditions are not interchangeable: some individuals with obesity retain relatively good insulin sensitivity, while insulin resistance can occur in people without obesity. Genetic susceptibility, physical activity, fat distribution, and other health conditions influence this variation. Consequently, body size alone cannot determine a person's metabolic status.

Understanding the relationship between obesity and insulin resistance is important because it helps explain why excess adiposity can contribute to abnormal glucose regulation. It also provides a biological basis for interventions that improve insulin sensitivity and reduce the risk of type 2 diabetes.

Insulin Action and Metabolic Regulation

Insulin coordinates the use and storage of nutrients after food intake. In skeletal muscle, it promotes glucose uptake and glycogen storage. In the liver, it helps regulate glucose production and storage. In adipose tissue, it suppresses the breakdown of stored fat, a process known as lipolysis.

When these tissues become insulin resistant, muscle glucose uptake decreases, suppression of hepatic glucose production becomes less effective, and fatty acid release from adipose tissue increases. These disturbances connect glucose metabolism with lipid metabolism and help explain why insulin resistance can affect several organs simultaneously.

Mechanisms Linking Obesity to Insulin Resistance

Adipose Tissue Dysfunction

Adipose tissue functions as both an energy reservoir and an endocrine organ. It releases hormones and other signaling molecules that influence metabolism throughout the body. During sustained energy excess, adipose tissue expands. In some individuals, this expansion is accompanied by enlarged adipocytes, cellular stress, altered blood supply, and changes in immune cell activity.

These changes can impair the ability of adipose tissue to store nutrients safely and respond normally to insulin. Visceral fat is particularly associated with adverse metabolic effects, although the mechanisms and their relative importance differ between individuals.

Ectopic Lipid Accumulation

When lipid storage and handling become disturbed, fat can accumulate in organs such as the liver and skeletal muscle. This is called ectopic lipid accumulation. Certain lipid intermediates, including diacylglycerols, can activate signaling pathways that interfere with insulin action.

In skeletal muscle, impaired insulin signaling reduces insulin-stimulated glucose transport. In the liver, altered signaling and increased delivery of fatty acids and glycerol can disturb glucose and lipid metabolism. These interactions demonstrate that the location and metabolic handling of fat are important alongside the total amount of adipose tissue.

Inflammation and Metabolic Stress

Obesity-associated adipose dysfunction often includes persistent, low-grade inflammation. Adipocytes and immune cells can release inflammatory mediators that alter cellular signaling and contribute to impaired insulin action.

However, the relationship is complex. Experimental research in mice has shown that adipose insulin resistance can precede macrophage accumulation and promote inflammation. Human tissue observations in the same study supported an association between impaired insulin signaling and inflammatory changes, but did not establish the same causal sequence in humans. These findings suggest that inflammation and insulin resistance can reinforce one another rather than follow a single universal pathway.

Increased Fatty Acid Release

Insulin normally restrains adipose tissue lipolysis. When this action becomes impaired, greater amounts of fatty acids and glycerol can reach the liver. These substrates contribute to hepatic triglyceride production and glucose production, potentially worsening metabolic dysfunction.

Thus, insulin resistance in adipose tissue can affect insulin-responsive organs elsewhere in the body. The resulting interaction between adipose tissue, muscle, and liver helps sustain abnormal nutrient handling.

Clinical Implications and Assessment

Insulin resistance and prediabetes frequently produce no obvious symptoms. They may coexist with hypertension, abnormal blood lipids, metabolic dysfunction-associated steatotic liver disease, and increased waist circumference. If pancreatic insulin secretion can no longer compensate adequately, blood glucose may rise and type 2 diabetes can develop.

Clinical assessment therefore considers medical history, body measurements, blood pressure, and relevant laboratory findings. Fasting plasma glucose, glycated hemoglobin, and oral glucose tolerance testing assess glucose regulation; they are not direct measurements of insulin resistance. Direct assessment of insulin resistance is primarily used in research rather than routine clinical practice.

Prevention and Management

Regular physical activity can improve insulin sensitivity and increase glucose uptake by skeletal muscle. These benefits make movement valuable even when changes in body weight are limited. For people with overweight or obesity, appropriate weight reduction can also improve insulin sensitivity. Management should use achievable goals and account for individual preferences and health circumstances.

Healthy eating, sufficient sleep, and sustained support are additional components of care. In adults at increased diabetes risk, the Diabetes Prevention Program demonstrated benefits from a lifestyle intervention that included a weight-loss goal of approximately 5–7%. Selected patients may also benefit from medication or metabolic surgery following clinical assessment. Treatment decisions depend on the presence of obesity, prediabetes, diabetes, and associated conditions.

Conclusion

Obesity and insulin resistance are connected through adipose tissue dysfunction, ectopic lipid accumulation, inflammatory processes, and altered nutrient metabolism. Their relationship varies between individuals and involves interactions among several organs. Effective care requires assessment of metabolic health alongside body weight, followed by individualized interventions that improve insulin sensitivity and reduce long-term disease risk.

Literatures:

1. Скворцова, Л. А., Махмудов, Д. Э., Усманов, Д. Р., Адылходжаев, А. А., & Хегай, Л. Н. (2019). АСПЕКТЫ СПОРТИВНОЙ АДАПТОЛОГИИ ПРИ ИЗУЧЕНИИ ВИТАМИНА D 25 (OH) D У ФУТБОЛИСТОВ. In От фундаментальных знаний к тонкому владению скальпелем (pp. 51-54).
2. Хегай, Л. Н., Скворцова, Л. А., Махмудов, Д. Э., Усманов, Д. Р., Муслимов, О. Я., & Адылходжаев, А. А. (2019). СОДЕРЖАНИЕ 25-ОН-ВИТАМИНА D У ФУТБОЛИСТОВ В УСЛОВИЯХ СОЛНЕЧНОЙ РАДИАЦИИ УЗБЕКИСТАНА. In Actual Questions and Innovations in Science 2 (pp. 160-161).
3. Mavlyanov, I., Parpiev, S., Sadikov, A., Kurganov, S., & Makhmudov, D. (2020). Relative features of the PPARA (Rs4253778), PPARGC1A (Rs8192678) and PPARG2 (Rs1801282) polymorphisms genes in athletes engaged in cyclic types of sports. European Journal of Molecular and Clinical Medicine, 7(2), 1860-1869.
4. Пирназаров, М. М., Садиков, А. А., Махмудов, Д. Э., & Хегай, Л. Н. (2021). Магнитно-резонансная томографическая визуализация повреждений коленных суставов. Re-health journal, (1 (9)), 152-159.

5. Махмудов, Д. Э., & Саитов, Т. А. (2023). Анализ персонализированного питания высококвалифицированных спортсменов управленческих видов спорта. Klinik va profilaktik tibbiyot jurnali, (3), 79.
6. Махмудов, Д. Э., & Саитов, Т. А. (2023). Анализ персонализированного питания высококвалифицированных спортсменов управленческих видов спорта. Klinik va profilaktik tibbiyot jurnali, (3), 79.
7. Таралева, Т. А., Махмудов, Д. Э., Атабаева, С. М., & Сагитов, Д. М. (2026). ВАРИАБЕЛЬНОСТЬ СЕРДЕЧНОГО РИТМА В ЦИКЛИЧЕСКИХ ВИДАХ СПОРТА: СОВРЕМЕННЫЕ ПОДХОДЫ К ОЦЕНКЕ ФУНКЦИОНАЛЬНОГО СОСТОЯНИЯ СПОРТСМЕНОВ. South Aral Region Medical Journal, 2(Maxsus son-2), 587-593.
8. Махмудов, Д., Абдуллаева, Х., Таралева, Т., Аблялимов, Р., & Парпиев, С. (2025). АНАЛИЗ ФУНКЦИОНАЛЬНЫХ ПОКАЗАТЕЛЕЙ ФУТБОЛИСТОВ НА ЭТАПЕ ОТБОРА К СОРЕВНОВАНИЯМ. Медицина и спорт, (3), 162-162.